

# BARBARA GRICE MEMORIAL SPAY AND NEUTER CLINIC

**Owner's Name** \_\_\_\_\_ **Cell Phone** \_\_\_\_\_ **Date** \_\_\_\_\_

**Owner's Street Address** \_\_\_\_\_ **City** \_\_\_\_\_ **County** \_\_\_\_\_ **State** \_\_\_\_\_ **Zip Code** \_\_\_\_\_

**Patient Name** \_\_\_\_\_ **Species Dog/Cat** \_\_\_\_\_ **Sex: Male/Female** \_\_\_\_\_

**Pet's Age: Years :** \_\_\_\_\_ **Months :** \_\_\_\_\_ **Pet's Breed** \_\_\_\_\_ **Pet's Color** \_\_\_\_\_

\_\_\_\_\_ cc Morphine 15mg/ml SQ IM      \_\_\_\_\_ cc Ketamine IV IM      \_\_\_\_\_ cc Midaz 5 mg/mL IV IM      \_\_\_\_\_ cc DetKetBut IM  
 \_\_\_\_\_ cc Ace 10mg/mL SQ IM      \_\_\_\_\_ cc Dilute Ace 1 mg/ml SQ IM      \_\_\_\_\_ cc 2% lidocaine block (testes)  
 \_\_\_\_\_ cc Meloxicam 5mg/mL SQ IM      \_\_\_\_\_ cc Dexium SP 4 mg/mL      \_\_\_\_\_ 0.1 cc 2% lidocaine-laryngeal folds  
 \_\_\_\_\_ cc mL LRS/ Saline IV SQ      \_\_\_\_\_ cc Atipamezole 5.0 mg/mL IM      \_\_\_\_\_ 2% Isoflurane + O2  
 \_\_\_\_\_ cc Cefovecin (Convenia) SQ      \_\_\_\_\_ cc Euthasol IV IC SQ Intrauterine      \_\_\_\_\_ SpO2, HR & RR Monitoring

Spay      **WEIGHT:** \_\_\_\_\_ (lbs)      S: BAR Abnormal      PreMed Time: \_\_\_\_\_  
 Neuter      O: Physical Exam  
 Already Sterilized      A: Surgical Candidate      Y      N  
 In Heat      P: Surgically Sterilize      Y      N  
 Pregnant # of Fetuses \_\_\_\_\_  
 Cryptorchid      T: (F) \_\_\_\_\_      P: (/min) \_\_\_\_\_      R: (/min) \_\_\_\_\_

**SPAY: Skin Incision:**      Ventral Midline      Flank - Right or Left (Ferals)  
 Ovarian Pedicles:      Instrument tie      Millers      Transfixation      Suture \_\_\_\_\_ P D O  
 Uterine Stump:      Millers      Transfixation      Suture \_\_\_\_\_ P D O  
 Abdominal Closure:      Cruciate      Simple Continuous      Suture \_\_\_\_\_ P D O  
 Subcutaneous Closure:      Simple Continuous      Simple Interrupted      Mattress      Suture \_\_\_\_\_ P D O  
 Intradermal Closure:      Simple Continuous      Simple Interrupted      Purse String Mattress (cat)      Suture \_\_\_\_\_ P D O  
 Incision Splash:      0.25% Bupivacaine (cat)      0.50% Bupivacaine/1% Epinephrine (9:1)      Phenylephrine (0.1 ml 10mg/ml to 10 ml saline) \_\_\_\_\_  
 Skin:      Surgical Glue      Staples \_\_\_\_\_      Ice Pack Incision Line Post Surgery

**NEUTER: Skin Incision:**      Pre-scrotal      Scrotal      **Technique:**      Closed Castration      Open Castration  
 Cord Ligation:      Instrument tie      Millers      Transfixation      Suture \_\_\_\_\_ P D O  
**Scrotal Closure:**      Single Suture Simple Subcutaneous Cruciate (dog scrotal)      Suture \_\_\_\_\_ P D O  
**Prescrotal Closure:**      2-layer Subcutaneous/Intradermal (dog pre-scrotal)      Suture \_\_\_\_\_ P D O  
 Incision Splash:      0.25% Bupivacaine (cat)      0.50% Bupivacaine/1% Epinephrine (9:1)      Phenylephrine (0.1 ml 10mg/ml to 10 ml saline) \_\_\_\_\_  
 Skin:      Surgical Glue      Staples \_\_\_\_\_      Ice Pack Scrotum Post Surgery

**ADDITIONAL CONCERNS ABOUT THIS PET**

Over / Underweight      Eye or Ear Concerns      Skin Abnormalities      Tapeworms      Dental Concerns      Fleas/Ticks  
 Other \_\_\_\_\_

**ADDITIONAL SERVICES**

Rabies      DA2PPV      Bordetella      FVRCP      FeLV      IV Fluids      Microchip      Dewormer \_\_\_\_\_  
 Fecal Parasite      Ear Cleaning      Nail Trim      Ear Tip      Capstar      Hernia Repair      **E-COLLAR**      Bitter Orange \_\_\_\_\_

**HW Test:**      -NAD (Neg)      + Pos Ag      **Fel/FIV Test**      Results:      FeLV: -Neg      +Pos      FIV: -Neg      +Pos

**Veterinarian:** \_\_\_\_\_